



NORMANHURST BOYS HIGH SCHOOL



MATHEMATICS ADVANCED (INCORPORATING EXTENSION 1) YEAR 11 COURSE



Topic summary and exercises:

(A) Algebraic Techniques & Coordinate Geometry

With references to



Name:

Initial version by H. Lam, 2018 (Algebraic Techniques), 2016 (Coordinate Geometry). Last updated February 3, 2025.
 Based on the work from the legacy syllabuses by R. Trenwith, 1995–2010, subsequently maintained by H. Lam, 2011–18.
 Additional editing by I. Ham and M. Ho, 2019–2020, and A. Sun in late 2020.
 Various corrections by students & members of the Mathematics Departments at North Sydney Boys and Normanhurst Boys High Schools.

Acknowledgements Pictograms in this document are a derivative of the work originally by Freepik at <http://www.flaticon.com>, used under  CC BY 2.0.

Symbols used

-  Beware! Heed warning.
-  Mathematics Advanced content.
-  Mathematics Extension 1 exclusive content.
-  Literacy: note new word/phrase.
-  Facts/formulae to memorise.
-  On the course Reference Sheet.
-  ICT usage
-  Enrichment content. Broaden your knowledge!
- $\mathbb{N}$ the set of natural numbers
- $\mathbb{Z}$ the set of integers
- $\mathbb{Q}$ the set of rational numbers
- $\mathbb{R}$ the set of real numbers
- $\forall$ for all

Syllabus outcomes addressed

MA11-1 uses algebraic and graphical techniques to solve, and where appropriate, compare alternative solutions to problems

MA11-2 uses the concepts of functions and relations to model, analyse and solve practical problems

Syllabus subtopics

MA-F1 Working with Functions

MA-F2 Graphing Techniques

Gentle reminder

- For a thorough understanding of the topic, *every* question in this handout is to be completed!
- Additional questions from *Cambridge Year 11 3 Unit* (Pender, Sadler, Shea, & Ward, 1999) will be completed at the discretion of your teacher.
- Remember to copy the question into your exercise book!

Learning intentions & outcomes

✓ Content/learning intentions

-
- ☐ 1.1 Perform binomial expansions:
- Square of a sum.
 - Square of a difference.
 - Difference of squares.
-
- ☐ 1.2 Can apply four standard methods of factorisation:
- Highest common factor.
 - Quadratic trinomial factorisations.
 - Difference of squares.
 - Grouping in pairs.
-
- ☐ 1.3 Manipulate complex algebraic expressions involving algebraic fractions.
- Add and subtract algebraic fractions, finding the LCD as appropriate.
 - Multiply and divide algebraic fractions, applying factorisation as appropriate.
 - Simplify compound fractions.
-
- ☐ 1.4 Solve linear equations and change the subject of a formula.
-
- ☐ 1.5 Solve quadratic equations using the quadratic formula and by completing the square  (ACMMM008)
- Also solve by factorisation.
-
- ☐ 1.6 Solve simultaneous equations (linear and quadratic) by substitution and elimination.
-
- ☐ 1.7 Use surd laws
- Express a surd in simplest form.
 - Simplify binomial expansions involving surds.
 - Simplify expressions involving $+$, $-$, $\times$, $\div$ of surds.
 - Rationalise the denominator.
-

Distance, Midpoint and Gradient in the Cartesian Plane

✓ Content/learning intentions

☐ 1.8 Distance formula between two points in the Cartesian Plane.

☐ 1.9 The midpoint formula.

☐ 1.10 Calculate the gradient of an interval.

Lines in the Cartesian Plane

✓ Content/learning intentions

☐ 1.11 Examine and use the relationship between the angle of inclination of a line or tangent θ with the positive x -axis, and the gradient m of that line or tangent, and establish that $\tan \theta = m$.

☐ 1.12 Understand and use the fact that parallel lines have the same gradient and that two lines with gradient m_1 and m_2 respectively are perpendicular if and only if $m_1 m_2 = -1$.

☐ 1.13 Test for the special quadrilaterals in the Cartesian plane using distance, midpoint and gradient calculations.

➤ A quadrilateral is a parallelogram is:

- opposites sides are equal length, or
- one pair of opposite sides are equal and parallel, or,
- the diagonals bisect each other.

➤ A quadrilateral is a rhombus if:

- all sides are equal length, or
- the diagonals bisect each other at right angles.

➤ A quadrilateral is a rectangle if:

- the diagonals are equal and bisect each other.
-

Equations of straight lines

✓ Content/learning intentions

- ☐ 1.14  Know and use the following forms of a straight line:
- Gradient-intercept form $y = mx + c$.
 - General form $ax + by + c = 0$.
-
- ☐ 1.15 Derive the equation of a straight line passing through a fixed point (x_1, y_1) and having a given gradient m using the formula $y - y_1 = m(x - x_1)$.
- *Point-gradient formula*
-
- ☐ 1.16 Derive the equation of a straight line passing through two points (x_1, y_1) and (x_2, y_2) by first calculating its gradient m using the formula $m = \frac{y_2 - y_1}{x_2 - x_1}$.
- *Two point formula*
-
- ☐ 1.17 Determine if three distinct lines are concurrent.
- Emphasise the need to resist finishing a proof with “ $1 = 1$ ” (from substituting both the x and y values into equations).
-
- ☐ 1.18 Find the equations of straight lines, including parallel and perpendicular lines, given sufficient information  (ACMMM004).
-
- ☐ 1.19 Solve a range of geometric problems using coordinate geometry concepts, including problems using pronumerals (rather than specific numeric values).
-

Contents

I	 Algebraic Techniques	7
1	Expansions, factorisations and algebraic fractions	8
1.1	Expansions and simplification	8
1.2	Factorisations	9
1.3	Algebraic Fractions	12
2	Equations	15
2.1	Linear equations and inequalities	15
2.2	Quadratic equations	16
2.2.1	Completing the square	16
2.2.2	Methods of solution	18
2.3	Simultaneous equations	20
2.3.1	Break-even analysis	22
3	Surd laws	27
3.1	Basic arithmetic of surds	27
3.2	Equality of surds	29
II	Coordinate Geometry	31
4	Points and intervals	32
4.1	Review of formulae	32
5	Gradient	34
5.1	Review of formulae	34
6	Equation of straight line	38
6.1	Review of formulae	38
	References	41

Part I

⌘ Algebraic Techniques

Section 1

Expansions, factorisations and algebraic fractions

1.1 Expansions and simplification



Learning Goal(s)



Knowledge
Stage 5 algebra



Skills
Factorisations and expansions



Understanding
The need to be able to perform expansions or factorisations to further simplify expressions

✓ **By the end of this section am I able to:**

- 1.1 Perform binomial expansions
- 1.2 Can apply four standard methods of factorisation
- 1.3 Manipulate complex algebraic expressions involving algebraic fractions

Only the most challenging questions are provided here. The rest are omitted for brevity and should be assumed knowledge.



Laws/Results

Square of a sum/difference

$$(a \pm b)^2 = \dots\dots\dots$$

Difference of squares

$$(a - b)(a + b) = \dots\dots\dots$$



Example 1



Expand and fully simplify $(a + b + c)^2$ by rewriting as $(a + (b + c))^2$.

Answer: $a^2 + b^2 + c^2 + 2ab + 2ac + 2bc$

1.2 Factorisations

Simply, to rewrite a sum of two or more terms into a of two or more terms.

Laws/Results

Four elementary methods of factorisation, which can be nested within each other:

Highest common factor

$$5a^2b + 10ab^2 = \dots\dots\dots$$

Difference of squares

$$a^2 - b^2 = \dots\dots\dots$$

Quadratic trinomial

$$a^2 - 4a - 77 = \dots\dots\dots$$

Grouping in pairs

$$3x - 3y + ay - ax = \dots\dots\dots$$

$$= \dots\dots\dots$$

Gentle reminder

For further revision, see the *A complete course on factorisations*, located at  [Lowe and Lam \(2010\)](#)

Exercises**1.** Simplify

$$(a) \quad 2a - 3b - a + 2b \qquad (b) \quad 5x^2 - 3x + x - x^2 + 2x \qquad (c) \quad a - (0.1)a$$

2. Simplify

$$\begin{array}{lll} (a) \quad 3a^4 \times 2a & (e) \quad (-2a^2b)^5 & (i) \quad x(2x+1)^3 \times x^4(2x+1) \\ (b) \quad 5a^3 \times 3a^2b^2 & (f) \quad 16x^8 \div 2x^2 & (j) \quad t \times \frac{1}{t} \\ (c) \quad (-3x) \times (-4x) & (g) \quad \frac{6x^3y^2z}{9x^2y^4z} & (k) \quad \sqrt{x} \times \sqrt{x} \\ (d) \quad (3a^2)^3 & (h) \quad (pq^3)^2 \times p^3q \div (pq)^3 & (l) \quad \text{reciprocal of } a + b \end{array}$$

3. (a) From $3x - 2y + 1$ take $x - y + 1$.(b) What must be added to $2m^2 - m + 1$ to give $5m^2 - 3m$?(c) How many times must p be subtracted from p^3 to give zero?(d) What is the product of x andi. half the reciprocal of x ?ii. the reciprocal of half x ?**4.** Expand, simplifying where necessary:

$$\begin{array}{lll} (a) \quad 3(2a - 5) & (h) \quad (w^3 + 2v^2)(w^3 - 2v^2) & (p) \quad (a - b)(a^2 + ab + b^2) \\ (b) \quad 3ab(2a + 5b) & (i) \quad 2(a - 3) - (1 - 2a) & (q) \quad (x - 2)(x^2 + 3x - 1) \\ (c) \quad -(1 - x) & (j) \quad 1 - (x - 1)^2 & (r) \quad (a - 1)(a - 2)^2 \\ (d) \quad (2x - 1)(x + 3) & (k) \quad (a + b)^2 - (a - b)(a + b) & (s) \quad (a - 2)^3 \\ (e) \quad (3a - 2)^2 & (l) \quad 3(v - 2)^2 & (t) \quad (2a + 1)^4 \\ (f) \quad (3q^2 + 2)^2 & (m) \quad (x - \sqrt{3})(x + \sqrt{3}) & (u) \quad \left(x + \frac{1}{x}\right)^2 \\ (g) \quad (1 - 2x)(1 + 2x) & (n) \quad a(a + 2)(3a - 4) & \\ & (o) \quad (a + b)(a - 2b + 1) & \end{array}$$

5. (a) Find an expansion for $(a + b + c)^2$ by first rewriting it as $[(a + b) + c]^2$. Try to guess the expansion for $(a + b + c + d)^2$.(b) A square has side length a . The length of the square is increased by b where $0 < b < a$, while the width decreased by b . Does the area increase or decrease? By how much?(c) In a right angled triangle, the length of the hypotenuse is $(x + 4)$ cm and the length of one of the short sides is $(x - 4)$ cm. Find an expression, in simplest form, for the length of the other short side.

6. Factorise *fully*:

- | | | |
|---------------------------|------------------------------------|---------------------------------|
| (a) $6a^2b^3 - 9ab^5$ | (j) $x^4 - 1$ | (r) $(a + b)^2 - c^2$ |
| (b) $a(b + 1) - b(b + 1)$ | (k) $4x^2 - 16y^2$ | (s) $a^2 - (b + c)^2$ |
| (c) $9a^2 - 16b^2$ | (l) $x^2 - 6x + 9$ | (t) $(a + b)^2 - (c - d)^2$ |
| (d) $-4x + 6$ | (m) $4x^2 + 12xy + 9y^2$ | (u) $x^2 - y^2 - x - y$ |
| (e) $x^2 - 3x - 54$ | (n) $\ell^4 - 4mn\ell^2 + 4m^2n^2$ | (v) $x^3 + 6x^2 + 9x$ |
| (f) $3x^2 - 8x - 3$ | (o) $6a^2 + 7ab - 3b^2$ | (w) $-x^2 + 2x + 15$ |
| (g) $12x^2 + x - 6$ | (p) $x^2 - 2\frac{1}{4}$ | (x) $x^4 - x^2 - 12$ |
| (h) $x^3 + x^2 + x + 1$ | (q) $\frac{a^2}{9} - \frac{4}{25}$ | (y) $a(a - b)^2 - ac^2$ |
| (i) $x^3 + x^2 - x - 1$ | | (z) $(x^2 - y^2)^2 - (x - y)^4$ |

7. Show that $(x + 1)^2 - (x - 1)^2 = 4x$ by

- (a) expanding and simplifying, and (b) factorising, without expanding.

8. Expand and factorise $1 - x(1 - x(1 - x))$.**9.** Factorise fully (without expanding):

- (a)
- $3(x + 1)^2 + 2x(x + 1)$
- (b)
- $x^4 - x^2(2x - 1)$
- (c)
- $x^2(x - 1)^3 - 2x(x - 1)^2$

10. If $f(n) = n(n + 1)(n + 2)$, express $f(n) + f(n + 1)$ in factored form. **Further exercises****(A) Ex 1B**

- Q3-7, 9

(A) Ex 1C

- Q1-7

(x1) Ex 1B

- Q7, 8, 9, 10

(x1) Ex 1C

- Q1-9

Answers

1. (a) $a - b$ (b) $4x^2$ (c) $\frac{9a}{10}$ or $0.9a$ 2. (a) $6a^5$ (b) $15a^5b^2$ (c) $12x^2$ (d) $27a^6$ (e) $-32a^{10}b^5$ (f) $8x^6$ (g) $\frac{2x}{3y^2}$ (h) p^2q^4 (i) $x^5(2x+1)^4$ (j) 1 (k) x (l) $\frac{1}{a+b}$ note: NOT $\frac{1}{a} + \frac{1}{b}$ 3. (a) $2x - y$ (b) $3m^2 - 2m - 1$ (c) p^2 times (d) i. $\frac{1}{2}$ ii. 2 4. (a) $6a - 15$ (b) $6a^2b + 15ab^2$ (c) $x - 1$ (d) $2x^2 + 5x - 3$ (e) $9a^2 - 12a + 4$ (f) $9q^4 + 12q^2 + 4$ (g) $1 - 4x^2$ (h) $w^6 - 4v^4$ (i) $4a - 7$ (j) $2x - x^2$ (k) $2b^2 + 2ab$ (l) $3v^2 - 12v + 12$ (m) $x^2 - 3$ (n) $3a^3 + 2a^2 - 8a$ (o) $a^2 - ab + a + b - 2b^2$ (p) $a^3 - b^3$ (q) $x^3 + x^2 - 7x + 2$ (r) $a^3 - 5a^2 + 8a - 4$ (s) $a^3 - 6a^2 + 12a - 8$ (t) $16a^4 + 32a^3 + 24a^2 + 8a + 1$ (u) $x^2 + 2 + \frac{1}{x^2}$ 5. (a) $a^2 + b^2 + c^2 + 2ab + 2ac + 2bc$; $a^2 + b^2 + c^2 + d^2 + 2ab + 2ac + 2ad + 2bc + 2bd + 2cd$ (b) decreases by b^2 (c) $4\sqrt{x}$ cm 6. (a) $3ab^3(2a - 3b^2)$ (b) $(b+1)(a-b)$ (c) $(3a-4b)(3a+4b)$ (d) $-2(2x-3)$ (e) $(x-9)(x+6)$ (f) $(3x+1)(x-3)$ (g) $(4x+3)(3x-2)$ (h) $(x+1)(x^2+1)$ (i) $(x+1)^2(x-1)$ (j) $(x-1)(x+1)(x^2+1)$ (k) $4(x-2y)(x+2y)$ (l) $(x-3)^2$ (m) $(2x+3y)^2$ (n) $(\ell^2 - 2mn)^2$ (o) $(3a-b)(2a+3b)$ (p) $(x - \frac{3}{2})(x + \frac{3}{2})$ (q) $(\frac{a}{3} - \frac{2}{5})(\frac{a}{3} + \frac{2}{5})$ (r) $(a+b+c)(a+b-c)$ (s) $(a+b+c)(a-b-c)$ (t) $(a+b+c+d)(a+b-c+d)$ (u) $(x+y)(x-y-1)$ (v) $x(x+3)^2$ (w) $-(x-5)(x+3)$ (x) $(x-2)(x+2)(x^2+3)$ (y) $a(a-b+c)(a-b-c)$ (z) $4xy(x-y)^2$ 7. "Show" question: 8. $(1-x)(1+x^2)$ 9. (a) $(x+1)(5x+3)$ (b) $x^2(x-1)^2$ (c) $x(x-1)^2(x+1)(x-2)$ 10. $(n+1)(n+2)(2n+3)$

1.3 Algebraic Fractions



Important note

- Obtain a prior to adding.
- Only multiply up by as much as required.



Example 2

Fully simplify:

(a) $\frac{1}{x-4} - \frac{1}{x}$

(b) $\frac{2}{x^2-x} - \frac{5}{x^2-1}$

Answer: (a) $\frac{4}{x(x-4)}$ (b) $\frac{2-3x}{x(x-1)(x+1)}$

Exercises

1. Simplify fully. Also state any values of pronumeral(s) for which the simplification is not valid.

(a) $\frac{4a+6}{4}$	(d) $\frac{x^2-2x-3}{x+1}$	(g) $\frac{4a-8}{12-6a}$	(j) $\frac{1-b^2}{b^3-1}$
(b) $\frac{4a+6}{6a+9}$	(e) $\frac{x+1}{x^2-2x-3}$	(h) $\frac{a^2-4}{a+2}$	(k) $\frac{x^4-5x^2+4}{x^2-x-2}$
(c) $\frac{x^2}{5x-x^2}$	(f) $\frac{1-x}{x-1}$	(i) $\frac{a^2+a-6}{a^2-3a+2}$	(l) $\frac{x^3+2x^2-4x-8}{x^2+4x+4}$

2. Write as a single fraction in simplest form.

(a) $\frac{a}{2} + \frac{2a}{3}$	(e) $\frac{4}{a} - \frac{3}{b}$	(i) $\frac{3}{a^3b} - \frac{2}{a^2b^4}$
(b) $\frac{m-1}{2} - \frac{2m-3}{5}$	(f) $\frac{1}{a^2} + \frac{2}{a}$	(j) $\frac{2}{x-2} + \frac{3}{x+1}$
(c) $x + \frac{1}{x}$	(g) $\frac{a+1}{2a} - \frac{a-2}{3a}$	(k) $\frac{2x}{x+3} - \frac{x-2}{x+1}$
(d) $a - \frac{a+b}{3}$	(h) $\frac{1}{(x-3)^2} + \frac{1}{x-3}$	(l) $\frac{1}{x^2+x} + \frac{1}{x^2-x}$
(m) $\frac{x}{(x-2)(x+2)} + \frac{x+1}{(x-2)(x-1)}$	(p) $\frac{1}{x^2-x-2} - \frac{1}{x^2+5x+4} - \frac{1}{x^2+2x-8}$	
(n) $\frac{2}{y+2} - \frac{1}{y+3} + \frac{5}{y-1}$	(q) $\frac{a^2}{a^2+3a+2} - \frac{2a}{a+2}$	
(o) $\frac{k}{k+1} + \frac{1}{k^2+3k+2}$		

3. Simplify fully:

(a) $\frac{4x^2}{3y^2} \times \frac{6y}{15x^4}$	(f) $\frac{9}{x^3+64} \div \frac{6}{x+4}$
(b) $\frac{3a^3}{7b^2} \div \frac{9a}{7b}$	(g) $\frac{x+y}{a-b} \times \frac{b-a}{y+x}$
(c) $\frac{2}{a} \div a$	(h) $\frac{a^2-a}{6a^3+6a^2} \div \frac{a^2-1}{8a}$
(d) $\frac{x+1}{2} \times \frac{4x}{(x+1)^2}$	(i) $\frac{a^2-2a-3}{a^2+3a} \times \frac{3a^2+18a+27}{a^2-9}$
(e) $\frac{1}{x+2} \times \frac{4x+8}{3}$	

4. x is the smallest of three consecutive integers.
- Find as a single fraction in simplest form, an expression for the sum of the reciprocals of these integers.
 - Three fractions are formed by dividing each of these integers by the integer following it. Find an expression, in simplest form, for the product of these fractions.
5. Find the reciprocal of $\frac{1}{a} + \frac{1}{b}$.
6. Simplify:
- $\frac{\frac{1}{m} + \frac{1}{n}}{m+n}$
 - $\frac{1}{1-\frac{m}{n}} + \frac{1}{1-\frac{n}{m}}$
 - $\frac{1-\frac{2}{t+1}}{t-\frac{2}{t+1}}$
7. (a) If $f(n) = n(n+1)(n+2)$, simplify $\frac{f(n)}{f(n+1)}$.
- (b) If $f(n) = \frac{n^2}{n-1}$, prove that $f\left(\frac{t}{t-1}\right) = f(t)$.

Answers

1. (a) $\frac{2a+3}{2}$ (b) $\frac{2}{3}$ [$a \neq -\frac{3}{2}$] (c) $\frac{x}{5-x}$ [$x \neq 0, 5$] (d) $x-3$ [$x \neq 1$] (e) $\frac{1}{x-3}$ [$x \neq -1, 3$] (f) -1 [$x \neq 1$] (g) $-\frac{2}{3}$ [$a \neq 2$]
 (h) $a-2$ [$a \neq -2$] (i) $\frac{a+3}{a-1}$ [$a \neq 1, 2$] (j) $-\frac{b+1}{b^2+b+1}$ [$b \neq 1$] (k) $(x+2)(x-1)$ [$x \neq -1, 2$] (l) $x-2$ [$x \neq -2$] 2. (a) $\frac{a}{6}$
 (b) $\frac{m+1}{10}$ (c) $\frac{x^2+1}{x}$ (d) $\frac{2a-b}{3}$ (e) $\frac{4b-3a}{ab}$ (f) $\frac{2a+1}{a^2}$ (g) $\frac{a+7}{6a}$ (h) $\frac{x-2}{(x-3)^2}$ (i) $\frac{3b^3-2a}{a^3b^4}$ (j) $\frac{5x-4}{(x-2)(x+1)}$ (k) $\frac{x^2+x+6}{(x+3)(x+1)}$ (l) $\frac{2}{x^2-1}$
 (m) $\frac{2(x^2+x+1)}{(x-2)(x+1)(x+2)}$ (n) $\frac{2(3y^2+14y+13)}{(y+2)(y+3)(y-1)}$ (o) $\frac{k+1}{k+2}$ (p) $\frac{-x+5}{(x-2)(x+1)(x+4)}$ (q) $-\frac{a}{a+1}$ 3. (a) $\frac{8}{15x^2y}$ (b) $\frac{a^2}{3b}$ (c) $\frac{2}{a^2}$ (d) $\frac{2x}{x+1}$ (e) $\frac{4}{3}$
 (f) $\frac{3}{2(x^2-4x+16)}$ (g) -1 (h) $\frac{4}{3(a+1)^2}$ (i) $\frac{3(a+1)}{a}$ 4. (a) $\frac{3x^2+6x+2}{x(x+1)(x+2)}$ (b) $\frac{x}{x+3}$ 5. $\frac{ab}{a+b}$ (NOT $a+b$!!!) 6. (a) $\frac{1}{mn}$ (b) 1 (c) $\frac{1}{t+2}$
 7. (a) $\frac{n}{n+3}$

Further exercises

(A) Ex 1D

- Q1-11 last column

(x1) Ex 1D

- Q2-13 last column

Section 2

Equations



Learning Goal(s)

Knowledge

Distinguishing the type of equation

Skills

Solve the equation using the appropriate technique

Understanding

The importance of being able to formulate and solve equations

By the end of this section am I able to:

- 1.4 Solve linear equations and change the subject of a formula.
- 1.5 Solve quadratic equations using the quadratic formula and by completing the square
- 1.6 Solve simultaneous equations (linear and quadratic) by substitution and elimination.

2.1 Linear equations and inequalities



Laws/Results

Two standard forms of linear equations:

1. - form

$$y = \dots\dots\dots$$

2. form

$$\dots\dots\dots = 0$$

where a is a whole number, and b and c are numbers.

Vertical/horizontal lines

- Vertical line: $\dots = a$

- Horizontal line: $\dots = b$

 Further exercises**(A) Ex 1E**

- Q1-3, 6-10

(x1) Ex 1E

- Q2-4, 7, 8, 11

(x1) Ex 5A

- Q3

2.2 Quadratic equations

2.2.1 Completing the square

**Example 3**

Complete the square for $x^2 + 4x + 1$ and rewrite as $(x - h)^2 + k$.

(See next page for the steps box)

**Steps**

1. Halve the coefficient of x , and then square it.
2. Complete the square:

Exercises

1. Complete the square:

(a) $x^2 + 6x + 1$	(d) $x^2 - 8x + 20$	(g) $a^2 - a$	(j) $x^2 + \frac{1}{2}x + 1$
(b) $x^2 - 2x - 5$	(e) $x^2 + x + 1$	(h) $4x^2 + 4x - 5$	(k) $x^2 + \frac{2}{3}x - 2$
(c) $a^2 + 4a$	(f) $x^2 - 3x - 3$	(i) $9x^2 - 24x + 14$	(l) $x^2 - \frac{3}{4}x$

2. Factorise by completing the square:

(a) $x^2 + 2x - 8$	(b) $a^2 + 2a - 1$	(c) $m^2 - 4m + 1$	(d) $4m^2 + 12m + 3$
--------------------	--------------------	--------------------	----------------------

3. Complete the square, then factorise:

(a) $x^2 + 2bx + c$	(b) $x^2 + \frac{b}{a}x + \frac{c}{a}$
---------------------	--

Answers

1. (a) $(x+3)^2 - 8$ (b) $(x-1)^2 - 6$ (c) $(a+2)^2 - 4$ (d) $(x-4)^2 + 4$ (e) $(x+\frac{1}{2})^2 + \frac{3}{4}$ (f) $(x-\frac{3}{2})^2 - \frac{21}{4}$ (g) $(a-\frac{1}{2})^2 - \frac{1}{4}$
 (h) $(2x+1)^2 - 6$ (i) $(3x-4)^2 - 2$ (j) $(x+\frac{1}{4})^2 + \frac{15}{16}$ (k) $(x+\frac{1}{3})^2 - \frac{19}{9}$ (l) $(x-\frac{3}{8})^2 - \frac{9}{64}$ 2. (a) $(x+4)(x-2)$ (b) $(a+1+\sqrt{2})(a+1-\sqrt{2})$
 (c) $(m-2+\sqrt{3})(m-2-\sqrt{3})$
 (d) $(2m+3-\sqrt{6})(2m+3+\sqrt{6})$ 3. (a) $(x+b+\sqrt{b^2-c})(x+b-\sqrt{b^2-c})$
 (b) $(x+\frac{b+\sqrt{b^2-4ac}}{2a})(x+\frac{b-\sqrt{b^2-4ac}}{2a})$

**Further exercises**

(A) Ex 1H
 • Q1-6

(x1) Ex 1H
 • Q1-6

2.2.2 Methods of solution

Laws/Results

Three methods of solution:

-
- the (rarely used as it is generalised into the method below)
-

Gentle reminder

Difference between an *equation* and *expression*

- An *equation* has an sign in the question. Commences with
- An *expression* does not have an sign in the question. Usually commences with , or

Important note

 When solving quadratic equations, always solve with as the subject!

**Example 4**

- (a) Fully factorise: $5x^2 + 34x - 7$
(b) Hence solve: $5x^2 + 34x - 7 = 0$

Answer: $\frac{1}{5}$ or -7 **Example 5**

Solve: $\frac{2}{a+3} + \frac{a+3}{2} = \frac{10}{3}$

Answer: $-\frac{7}{3}$ or 3 **Further exercises**

- Ⓐ Ex 1F
• Q2-11

- ⓧ Ex 1F
• Q3-13

2.3 Simultaneous equations



Important note

Three methods of solution:

1.
 - The number of solutions to a pair of equations is geometrically, the number of of two corresponding graphs.
2.
3.



Example 6

Solve:
$$\begin{cases} 3x - 2y = 29 \\ 4x + y = 24 \end{cases}$$

Answer: $x = 7, y = -4$



Example 7

Solve:
$$\begin{cases} y = x^2 \\ y = x + 2 \end{cases}$$

Answer: $x = 2, y = 4$ or $x = -1, y = 1$



Important note

After substituting, what type of equation shows up?

**Example 8**

Solve:
$$\begin{cases} x^2 + y^2 = 53 \\ x^2 - y^2 = 45 \end{cases}$$

Answer: $x = 7, y = \pm 2$ or $x = -7, y = \pm 2$

2.3.1 Break-even analysis

 One application of simultaneous equations is to find the break-even point in manufacturing and sales.

Fill in the spaces

- : A formula for the cost to produce a given number of items
- : the sale price multiplied by the number of units sold

Definition 1

..... - is when the cost function and revenue function



Example 9

(Fitzpatrick & Aus, 2018, p.125) Georgia decides to turn her jewellery-making hobby into a small business. She spends \$120 on equipment and estimates that it costs her \$8 to manufacture each item. She plans to sell the items for \$20 each.


- If C is her total cost and R her total revenue, in dollars, set up the cost and revenue functions for the sale of x items.
- Determine her break-even point.
- Determine her profit if she sells:
 - 6 items
 - 50 items

 Example 10

[2020 Adv HSC Q11] There are two tanks on a property, Tank *A* and Tank *B*. Initially, Tank *A* holds 1 000 litres of water and Tank *B* is empty.

- (a) Tank *A* begins to lose water at a constant rate of 20 litres per minute. 1
The volume of water in Tank *A* is modelled by $V = 1000 - 20t$ where V is the volume in litres and t is the time in minutes from when the tank begins to lose water.

On the grid below, draw the graph of this model and label it as Tank *A*.



**Example 11**

[2020 Adv HSC Q11] (*continued from previous page*)

- (b) Tank B remains empty until $t = 15$ when water is added to it at a constant rate of 30 litres per minute. **2**

By drawing a line on the grid on the previous page, or otherwise, find the value of t when the two tanks contain the same volume of water.

- (c) Using the graphs drawn, or otherwise, find the value of t (where $t > 0$) when the total volume of water in the two tanks is 1000 litres. **1**

Exercises

Source: [Fitzpatrick and Aus \(2018, Ex 5.5\)](#), unless otherwise stated.


1. For each set of cost and revenue functions (a)-(d), find:
 - i. the break-even point
 - ii. the revenue at the break-even point
 - iii. the profit function, P
 - (a) $C = 5x + 200, R = 15x$
 - (b) $C = 0.5x + 100, R = 1.5x$
 - (c) $C = 15x + 3000, R = 45x$
 - (d) $C = 0.3x + 5000, R = 1.1x$
2. Julian buys a coffee cart for \$15 000. His repayments work out to be \$80 per day for the first year. He calculates that it will cost him \$2 per cup of coffee for the ingredients. He sells coffee at \$4 per cup and he sells x cups of coffee per day.
 - (a) Write the cost function C and revenue function R .
 - (b) What is his break-even point?
 - (c) Write the profit function P .
 - (d) What is his profit if he sells 100 cups per day?
3. Maya runs a market stall at the weekends, selling paintings. It costs her \$90 per day for the site. It costs her on average \$4 per painting that she sells and she sells them for an average price of \$10. If she sells x articles each day, find:
 - (a) the cost function C and revenue function R .
 - (b) her break-even point
 - (c) the profit she will make if she sells 40 paintings in a day.
 - (d) One weekend, the weather is fine on the Saturday and rainy on the Sunday. On Saturday Maya sells 30 paintings, but on Sunday she only sells 10 paintings. What profit (or loss) does she make for the weekend?

4. [2020 CSSA Adv Trial Q14] Michael has a small manufacturing business.

The cost of manufacturing is given by the equation $C = 50x + 200$ and the income earned is given by the equation $I = 100x$, where x is the number of items that the business has manufactured.

- (a) Graph each of the two equations on the grid below.

2



- (b) How many items need to be manufactured for the business to break even?

1

Answers

1. (a) i. $x = 20$ ii. $R = \$300$ iii. $P = 10x - 200$ (b) i. $x = 100$ ii. $R = \$150$ iii. $P = x - 100$ (c) i. $x = 100$ ii. $R = \$4500$ iii. $P = 30x - 3000$ (d) i. $x = 6250$ ii. $R = \$6875$ iii. $P = 0.8x - 5000$ 2. (a) $C = 2x + 80$, $R = 4x$ (b) $x = 40$ (c) $P = 2x - 80$ 3. (a) $C = 4x + 90$, $R = 10x$ (b) $x = 15$ (c) $P = 6x - 90$, $x = 40$, $P = \$150$ (d) Saturday $x = 30$, $P = \$90$ / Sunday $x = 10$, $P = -\$30$ / Profit for the weekend: $\$60$ 4. 4 items

Further exercises

(A) Ex 1G
• Q3-5

(x1) Ex 1G
• Q3-5

Section 3

Surd laws



Learning Goal(s)

Knowledge

Simplifying surds

Skills

Rewriting surds in lowest base and rationalising

Understanding

The difference between *simplifying* and *rationalising* surds

By the end of this section am I able to:

1.7 Use surd laws

3.1 Basic arithmetic of surds



Theorem 1

Fully simplify surds by finding their factor.
.....



Example 12

Fully simplify: $\sqrt{72} - \sqrt{50} + \sqrt{12}$

Answer: $\sqrt{2} + 2\sqrt{3}$



Example 13

Expand and fully simplify: $(\sqrt{15} - \sqrt{6})^2$

Answer: $21 - 6\sqrt{10}$

 Theorem 2

Rewrite surds with a denominator by multiplying numerator and denominator by the of the denominator.

 Example 14

Fully simplify: $\frac{1}{2\sqrt{3} - 3\sqrt{2}}$.

Answer: $-\frac{2\sqrt{3}+3\sqrt{2}}{6}$

3.2 Equality of surds

Theorem 3

Two surds $a + b\sqrt{c}$ and $x + y\sqrt{c}$ are equal iff and

Example 15

Find the value of a and b if $\frac{2}{\sqrt{5}+1} = a + b\sqrt{5}$.

Answer: $a = -\frac{1}{2}, b = \frac{1}{2}$

Exercises

Acknowledgement: Portions taken from [Grove \(2010, Ex 2.23\)](#)

1. Find the values of a and b if

$$(a) \quad \frac{3}{2\sqrt{5}} = \frac{\sqrt{a}}{b}$$

$$(c) \quad \frac{2\sqrt{7}}{\sqrt{7}-4} = a + b\sqrt{7}$$

$$(b) \quad \frac{\sqrt{3}}{4\sqrt{2}} = \frac{a\sqrt{6}}{b}$$

$$(d) \quad \frac{\sqrt{2}+3}{\sqrt{2}-1} = a + \sqrt{b}$$

2. Show that $\frac{\sqrt{2}-1}{\sqrt{2}+1} + \frac{4}{\sqrt{2}}$ is rational.

3. If $x = \sqrt{3} + 2$, fully simplify:

$$(a) \quad x + \frac{1}{x}$$

$$(b) \quad \left(x + \frac{1}{x}\right)^2$$

$$(c) \quad x^2 + \frac{1}{x^2}$$

4. If $2 + \frac{1}{x} = \sqrt{3}$ where $x \neq 0$, find the exact value of x (with a rational denominator).

Answers

1. (a) $a = 45, b = 10$ (b) $a = 1, b = 8$ (c) $a = -\frac{14}{9}, b = -\frac{8}{9}$ (d) $a = 5, b = 32$ 2. Show 3. (a) 4 (b) 16 (c) 14 4. $-(\sqrt{3} + 2)$

Further exercises

(A) Ex 2C

- Q1-8 last column

(A) Ex 2D

- Q1-11 last 2 columns

(A) Ex 2E

- last 2 columns

(x1) Ex 2C

- Q1-8 last column

(x1) Ex 2D

- Q1-13 last 2 columns

(x1) Ex 2E

- last 2 columns

Part II

Coordinate Geometry

Section 4

Points and intervals



Learning Goal(s)



Knowledge

The distance and midpoint formula in the Cartesian plane



Skills

Finding the distance and midpoint of an interval



Understanding

The derivation of the distance and midpoint formulas

✓ By the end of this section am I able to:

- 1.8 Distance formula between two points in the Cartesian Plane
- 1.9 The midpoint formula

4.1 Review of formulae



Laws/Results



The distance formula to calculate the distance between two points (x_1, y_1) and (x_2, y_2) :

$$d = \dots\dots\dots$$

Derivation of formula:



Laws/Results



The midpoint formula to calculate the midpoint between two points (x_1, y_1) and (x_2, y_2) :

$$M(x, y) = \dots\dots\dots$$

Derivation of formula:

**Example 16**

The interval joining $A(3, -7)$ and $B(-6, 2)$ is a diameter of a circle. Find the centre and radius of the circle.

Answer: $C(-\frac{3}{2}, -\frac{5}{2})$, $r = \frac{9}{2}\sqrt{2}$

**Further exercises****② Ex 6A**

- Q1-2 last column
- Q3-10
- Q16-17

Section 5

Gradient



Learning Goal(s)



Knowledge

The relation between parallel and perpendicular lines



Skills

Determine the type of quadrilateral by examining the gradient of its sides and diagonals



Understanding

The relation between the gradient and the angle of inclination

✓ By the end of this section am I able to:

- 1.10 Calculate the gradient of an interval
- 1.11 Examine and use the relationship between the angle of inclination of a line or tangent θ with the positive x -axis, and the gradient m of that line or tangent, and establish that $\tan \theta = m$
- 1.12 Understand and use the fact that parallel lines have the same gradient and that two lines with gradient m_1 and m_2 respectively are perpendicular if and only if $m_1 m_2 = -1$
- 1.13 Test for the special quadrilaterals in the Cartesian plane using distance, midpoint and gradient calculations

5.1 Review of formulae



Laws/Results



The gradient formula to calculate the gradient between two points (x_1, y_1) and (x_2, y_2) :

$$m = \dots\dots\dots$$

Derivation of formula:



Laws/Results

The angle of inclination θ of the line and the positive direction of the x axis:

$$m = \dots\dots\dots$$

Derivation of formula:

 Laws/Results

Parallel lines have the gradient.

 Laws/Results

The gradient of two perpendicular lines are
..... :
..... = or

 Example 17

Show that the points $A(-2, 5)$, $B(1, 3)$ and $C(7, -1)$ are collinear

 Example 18

[Ex 5B Q16] Answer the following for the points $W(2, 3)$, $X(-7, 5)$, $Y(-1, -3)$ and $Z(5, -1)$:


- (a) Show that WZ is parallel to XY .
- (b) Find the lengths WZ and XY . Hence deduce the type of quadrilateral $WXYZ$.
- (c) Show that that the diagonals WY and XZ are perpendicular.

**Example 19****[Ex 5B Q18]**

- (a) $A(1, 4)$, $B(5, 0)$ and $C(9, 8)$ form the vertices of a triangle. Find the coordinates of P and Q if they divide the sides AB and AC respectively in the ratio $1 : 3$.
- (b) Show that PQ is parallel to BC and is one quarter of its length.

**Example 20**

The points A , B and C have coordinates $(1, 0)$, $(0, 8)$ and $(7, 4)$, and the angle between AC and the x axis is θ .



- (a) Find the gradient of the line AC and hence determine θ to the nearest degree. **2**
- (b) Find the equation of AC . **2**
- (c) Find the coordinates of D , the midpoint of AC . **2**
- (d) Show that $AC \perp BD$. **1**
- (e) What type of triangle is ABC ? Show full reasoning. **2**
- (f) Find the area of this triangle. **2**
- (g) Write down the coordinates of the point E such that $ABCE$ is a rhombus. **2**

Further exercises
② Ex 6B

- Q7-21 odd #

ⓧ1 Ex 5B

- Q7-21 odd #

Section 6

Equation of straight line



Learning Goal(s)

Knowledge

The different forms of straight lines

Skills

Finding the equation of straight lines

Understanding

Applying the appropriate form of a straight line to find its equation

By the end of this section am I able to:

- 1.14 Know and use the following forms of a straight line: gradient-intercept form $y = mx + c$, and general form $ax + by + c = 0$
- 1.15 Derive the equation of a straight line passing through two points (x_1, y_1) and (x_2, y_2) by first calculating its gradient m using the formula $m = \frac{y_2 - y_1}{x_2 - x_1}$
- 1.16 Derive the equation of a straight line passing through two points (x_1, y_1) and (x_2, y_2) by first calculating its gradient m using the formula $m = \frac{y_2 - y_1}{x_2 - x_1}$
- 1.17 Determine if three distinct lines are concurrent
- 1.18 Find the equations of straight lines, including parallel and perpendicular lines, given sufficient information

6.1 Review of formulae



Laws/Results



The gradient-intercept form of the equation of a line:

.....

where

• — • —



Laws/Results



The general form of the equation of a line:

.....

 Laws/Results

 The **point-gradient formula** to find the equation of a straight line through (x_1, y_1) with a given gradient m :

.....

Derivation of formula:

 Important note

No new theory is in this section. However, expect slightly more difficult problems within textbook exercises.

 Example 21

[Ex 5D Q6] Given the points $A(1, -2)$ and $B(-3, 4)$, find in general form the equation of:

- (a) the line AB ,
- (b) the line through A perpendicular to AB .

**Example 22****[Ex 5D Q12]**

- (a) On a number plane, plot the points $A(4, 3)$, $B(0, -3)$ and $C(4, 0)$.
- (b) Find the equation of BC .
- (c) Explain why $OACB$ is a parallelogram.
- (d) Find the area of $OACB$ and the length of the diagonal AB .

**Further exercises****② Ex 6C**

- Q5, 10-16

② Ex 6D

- Q5
- Q7 last 3 columns
- Q14, 18-20

ⓧ Ex 5C

- Q6, 10-14

ⓧ Ex 5D

- Q4, 7-10, 13-18

References

- Fitzpatrick, J. B., & Aus, B. (2018). *New Senior Mathematics – Advanced Course for Years 11 & 12* (3rd ed.). Pearson Education.
- Grove, M. (2010). *Maths in focus: mathematics extension preliminary course* (E. Bron, Ed.). McGraw-Hill Australia Pty Ltd.
- Lowe, R., & Lam, H. (2010). *A complete course on factorisation*. Normanhurst Boys High School Edmodo site. (Compiled by R. Lowe. Typeset in 2010 and edited in 2013 by H. Lam)
- Pender, W., Sadler, D., Shea, J., & Ward, D. (1999). *Cambridge Mathematics 3 Unit Year 11* (1st ed.). Cambridge University Press.
- Pender, W., Sadler, D., Shea, J., & Ward, D. (2009). *Cambridge Mathematics 2 Unit Year 11* (2nd ed.). Cambridge University Press.
- Trenwith, R., & Lam, H. (2018). *Algebraic Techniques, Functions and Further Work with Functions: Exercises for Topics 1, 2 and 3 (Mathematics Advanced incorporating Mathematics Extension 1, Year 11 Course)*. Normanhurst Boys High School Edmodo groups. (First compiled by R. Trenwith. Edited 2011, 2013, 2015, 2018 by H. Lam)